Online Immunization Reporting and Viewing

Every time your child gets immunized you need to report it to the Durham Region Health Department.

You can view or report immunizations online using the Immunization Connect Ontario (ICON) Tool. ICON allows you to report and view your child's immunizations online any time. Please also attach a photo of the record.




To report online, you will need the student's health card

- 1. Visit the website: durham.ca/immunize or Scan QR code
- 2. Launch ICON
- 3. Follow the instructions.
- 4. See the back of this page for reporting tool













Immunization Connect Ontario (ICON) Reporting Tool

If entering vaccine information online using ICON, please type the vaccine type as is written below.

2 months of age (Three immunizations due)		
	DTaP-IPV-Hib Pneu-C Rot-5	Diphtheria, tetanus, pertussis, polio, Haemophilus influenzae type b vaccine Pneumococcal conjugate 13 vaccine (Ex. Prevnar 13 ®) Rotavirus vaccine (Ex. RotaTeq ®)
4 n	nonths of age (Th	ree immunizations due)
	DTaP-IPV-Hib	Diphtheria, tetanus, pertussis, polio, Haemophilus influenzae type b vaccine
	Pneu-C Rot-5	Pneumococcal conjugate 13 vaccine (Ex. Prevnar 13 ®) Rotavirus vaccine (Ex. RotaTeq ®)
6 n	nonths of age (Tw	vo immunizations due)
	DTaP-IPV-Hib Rot-5	Diphtheria, tetanus, pertussis, polio, Haemophilus influenzae type b vaccine Rotavirus vaccine (Ex. RotaTeq ®)
12 months of age (Three immunizations due)		
	MMR Men-C-C Pneu-C	Measles, mumps and rubella vaccine (Ex. Priorix ®, M-M-R II ®) Meningococcal conjugate type C vaccine (Ex. Menjugate ®, Meningitec ®, NeisVac-C ®) Pneumococcal conjugate 13 vaccine (Ex. Prevnar 13 ®)
	oses of MMR vac gardless of previ	ccine, Men-C-C vaccine and Pneu-C-13 vaccine are needed on or after age one ous doses.
15	months of age (C	One immunization due)
	Var	Varicella (chickenpox) vaccine
18	months of age (C	One immunization due)
	DTaP-IPV-Hib	Diphtheria, tetanus, pertussis, polio, Haemophilus influenzae type b vaccine
4 y	ears of age (Two	immunizations due)
	Tdap-IPV	Tetanus, diphtheria, pertussis and polio vaccine (Ex. Adacel-Polio ®)
	MMRV	Measles, mumps, rubella and varicella vaccine (Ex. Priorix-Tetra ®, ProQuad ®)
* A	dose of Tdap-IF	V vaccine is needed on or after 4 years of age regardless of previous doses.
Gra	ade 7 vaccines (F	Report online ONLY if it was received outside of Durham Region Health, i.e. school or catchup)
	Men-C-ACYW13	35 -Meningococcal conjugate type ACYW135 vaccine (Ex. Menveo ®, Menactra ®, Nimenrix ®
	НВ	Hepatitis B (Ex. Recombivax ®, Engerix B ®)
	HPV9	Human papillomavirus (Ex. Gardasil 9 ®)
14-	- 16 year old boos	ster (10 year booster) Most common missing vaccine for high school students
	Tdap	Tetanus, diphtheria, pertussis vaccine (Ex. Adacel ®, Boostrix ®)